

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
<p>The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to the Department of Defense, Executive Service Directorate (0704-0188). Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.</p> <p>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ORGANIZATION.</p>					
1. REPORT DATE (DD-MM-YYYY) 28/06/2018		2. REPORT TYPE Poster		3. DATES COVERED (From - To) 06/28/2018-06/29/2018	
4. TITLE AND SUBTITLE Incorporation of Mindfulness Exercises to Reduce Anxiety and Pain during Urodynamic Testing: A Randomized Controlled Pilot Study				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Uberoi, Pansy Maj				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) 59th Clinical Research Division 1100 Willford Hall Loop, Bldg 4430 JBASA-Lackland, TX 78236-9908 210-292-7141				8. PERFORMING ORGANIZATION REPORT NUMBER 17668	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) 59th Clinical Research Division 1100 Willford Hall Loop, Bldg 4430 JBASA-Lackland, TX 78236-9908 210-292-7141				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release. Distribution is unlimited.					
13. SUPPLEMENTARY NOTES SURF Meeting, San Antonio, TX, June 28-29, 2018					
14. ABSTRACT Incorporation of Mindfulness Exercises to Reduce Anxiety and Pain during Urodynamic Testing: A Randomized Controlled Pilot Study Pansy Uberoi MD, MPH, Anna Smitherman PhD, Forrest Jellison MD San Antonio, TX. Presentation to be made by Dr. Uberoi Objectives: Mindfulness exercises have shown improvement in emotional coping and reduction of anxiety, leading to their incorporation into treatment for acute and chronic pain conditions. The present trial compares the impact of mindfulness exercises as they pertain to anxiety and pain levels among patients undergoing invasive in-office urodynamics (UDS). Methods: Twenty-seven patients were randomized to either psychologist led mindfulness exercises or an empty quiet room prior to undergoing UDS. The personnel performing UDS were blinded to the intervention. The primary outcome, reduction of anxiety after UDS, was measured by the					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON Clarice Longoria
a. REPORT	b. ABSTRACT	c. THIS PAGE			19b. TELEPHONE NUMBER (Include area code) 210-268-2201



Incorporation of Mindfulness Exercises to Reduce Anxiety and Pain During Urodynamic

Testing: a Randomized Controlled Pilot Study

Pansy Uberoi MD, MPH; Forrest Jellison MD; Anna Smitherman PhD

San Antonio Uniformed Services Health Education Consortium, San Antonio, TX




Objective

Mindfulness exercises have shown improvement in emotional coping and reduction of anxiety, leading to incorporation in treatment plans for chronic and acute pain conditions^{1,2}. The present randomized pilot study compares the impact of mindfulness among patients undergoing invasive in-office urodynamic studies^{3,4} with primary outcome of anxiety and secondarily pain, fear and embarrassment.

Methods

Randomized, controlled, pilot study




Data collected

- Visual Analog Scale for Pain (VAS)
- State-Trait Anxiety Inventory (STAI-6)
- Participant Questionnaire

Primary endpoint – anxiety measured by STAI-6




Secondary endpoints – pain, fear, embarrassment



Results

Demographics	Mindfulness	Control	p-value
Age [mean and (sd)]	59.23 (10.83)	57.46 (14.85)	0.73
Male (n)	8% (1)	0% (0)	
Female (n)	92% (12)	100% (13)	0.23
Race (n)			
—White	38% (5)	54% (7)	
—Black	31% (4)	15% (2)	
—Hispanic	23% (3)	23% (3)	
—Other	8% (1)	8% (1)	0.80
Perceived Health* [mean and (sd)]	2.54 (0.66)	1.85 (0.69)	0.02

Symptoms	Mindfulness (n)	Control (n)	p-value
Leakage/Incontinence	69% (9)	77% (10)	0.66
Obstruction/Blockage	31% (4)	15% (2)	0.35
Urgency	77% (10)	62% (8)	0.39
Frequency	54% (7)	62% (8)	0.69



Emotions	Mindfulness	Control	p-value
Anxious	1.00 (1.08)	1.31 (1.11)	0.45
Afraid	0.08 (0.28)	0.85 (1.21)	0.05
Embarrassed	0.54 (0.88)	1.23 (1.09)	0.07
Uncomfortable	1.25 (1.22)	1.23 (1.36)	0.80
Pain	0.92 (1.04)	1.31 (1.38)	0.35
Bladder Pain	1.39 (1.12)	0.62 (0.18)	0.05
Rectal Pain	0.92 (0.49)	1.08 (1.04)	0.80
Worry about Radiation	0.08 (0.28)	0.23 (0.60)	0.52
Worry about UTI	0.15 (0.56)	0.92 (1.32)	0.04
Worry about Results	0.54 (0.66)	0.62 (0.77)	0.86

Discussion

Urodynamics is associated with anxiety, embarrassment and discomfort³. Mindfulness exercises have been reported to improve emotional coping in acute and chronic conditions^{1,2}.

In this pilot randomized control trial, mindfulness exercises prior to UDS did not show an improvement in the primary outcomes anxiety as measured by the STAI-6. Secondary outcomes of fear, embarrassment, and worry about UTI were noted to be lower in the mindfulness group with statistical significance or trend toward it. As hypothesized, there was no significant difference in pain levels on VAS. Bladder pain was significantly worse in the mindfulness group. When taken in aggregate, anxiety levels decreased significantly for all patients.

We suspect a confounding effect may have been present in the control group as several subjects self-reported performing their own "relaxation techniques" in the quiet, empty room.

Summary

1. Mindfulness exercises prior to UDS did not result in decreased anxiety or pain.
2. Fear and embarrassment were significantly lower or trending in the mindfulness group.
3. Future study with a better designed control that simulates the clinical environment with adequate power may impact anxiety and pain during UDS.

References

1. Erisman SM, Roemer L. A Preliminary investigation of the effects of experimentally-induced mindfulness on emotional responding to film clips. *Emotion*, 10(1):72-82, 2010
2. Kanter G, Komosu YM. Mindfulness-based stress reduction as a novel treatment for interstitial cystitis/bladder pain syndrome: a randomized controlled trial. *Int Urogynecol J*, 27:1705, 2016
3. Suskind AM, Clemens JQ. Patient perceptions of physical and emotional discomfort related to urodynamic testing: a questionnaire-based study in men and women with and without neurologic conditions. *Urology*, 85(3):547-51, 2015
4. Solomon ER, Ridgway B. Interventions to decrease pain and anxiety in patients undergoing urodynamic testing: a randomized controlled trial. *NeuroUrol Urodyn*, 35:975-9, 2016

Disclaimer: The views expressed are those of the authors and do not reflect the official views or policy of the Department of Defense or its Components. The voluntary, fully informed consent of the subjects used in this research was obtained as required by 32CFR219 and DODI

3216.02_AFI 40-204

**Incorporation of Mindfulness Exercises to Reduce Anxiety and Pain during
Urodynamic Testing: A Randomized Controlled Pilot Study**

Pansy Uberoi MD, MPH, Anna Smitherman PhD, Forrest Jellison MD
San Antonio, TX. Presentation to be made by Dr. Uberoi

Objectives: Mindfulness exercises have shown improvement in emotional coping and reduction of anxiety, leading to their incorporation into treatment for acute and chronic pain conditions. The present trial compares the impact of mindfulness exercises as they pertain to anxiety and pain levels among patients undergoing invasive in-office urodynamics (UDS).

Methods: Twenty-seven patients were randomized to either psychologist led mindfulness exercises or an empty quiet room prior to undergoing UDS. The personnel performing UDS were blinded to the intervention. The primary outcome, reduction of anxiety after UDS, was measured by the state-trait anxiety inventory (STAI-6). STAI-6 consists of questions regarding positive aspects of mood and negative aspects of mood. Secondary outcome of pain was measured by the visual analog scale (VAS). Participants completed questionnaires at baseline, after control/study intervention, and immediately after UDS. Wilcoxon-Mann-Whitney test was performed for difference between anxiety and pain between groups. Secondary analyses were performed with Wilcoxon sign rank test to measure differences in changes within each group.

Results: Demographics, past medical history and presenting symptoms were similar between the groups. There was no statistically significant difference in anxiety in the mindfulness or control group at baseline ($p=0.42$), after intervention ($p=0.15$), or after UDS ($p=0.13$). Pain was not statistically different between groups at baseline ($p=0.17$), after intervention ($p=0.11$), after UDS ($p=0.38$). Secondary outcomes of fear ($p=0.05$) and embarrassment ($p=0.07$) were lower in the mindfulness group. When taken in aggregate, anxiety levels, as measured by STAI-6 were significantly different for all participants mean 12.00 at baseline and 12.46 after UDS ($p=0.05$).

Conclusions: This pilot study does not reflect reduction in patients' anxiety and pain during urodynamics procedure. Fear and embarrassment were noted to be lower in the mindfulness group. Decrease in anxiety was noted in all patients. We suspect confounding effect may have been present in the control group based on self-report of 'relaxation techniques' used by control patients in the quiet, empty room.

Source of Funding: None

Disclaimer: The views expressed are those of the authors and do not reflect the official views or policy of the Department of Defense or its Components. The voluntary, fully informed consent of the subjects used in this research was obtained as required by 32CFR219 and DODI 3216.02_AFI 40-204